A STATE OF THE STA		532 Rec'd PCT/PTC 27	SEP 2000			
FORM PTC-1390 (REV 11-98)	U.S DEPARTMENT OF COMMERCE FATENT AND T	TRADEMARK OFFICE ATTORNEY'S DOCKET NUMBER				
	ETTER TO THE UNITED	STATES				
DESIGNATED/E	ELECTED OFFICE (DO/		TR 1.51			
CONCERNING A	A FILING UNDER 35 U.S	S.C. 371 09/647329				
INTERNATIONAL APPLICATION		G DATE PRIORITY DATE CLAIMED				
PCT/GB99/00968 TITLE OF INVENTION	March 26, 1999	March 27, 1998				
POLYM		THMA TO A LOCUS ON CHROMOSOME 2				
APPLICANT(S) FOR DO/EO/US SUMIT BHA	WILLIAM OSMOND CHARLES M TTACHARYYA; NICHOLAS LEA	ICHAEL COOKSON; MIRIAM FLEUR MOFFATT;	;			
		(DO/EO/US) the following items and other information:				
1. X This is a FIRST submiss	ion of items concerning a filing under 3	35 U.S.C. 371.				
	BSEQUENT submission of items con		İ			
examination until the exp	iration of the applicable time limit set	(35 U.S.C. 371(f)) at any time rather than delay in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1) is made by the 19th month from the earliest claimed priority	date			
_ []	onal Application as filed (35 U.S.C	-	Siste.			
a. is transmitted		smitted by the International Bureau).	l			
· · · · · · · · · · · · · · · · · · ·	mitted by the International Bureau.					
		United States Receiving Office (RO/US).				
	ternational Application into English	on under PCT Article 19 (35 U.S.C. 371(c)(3))				
		insmitted by the International Bureau).				
b. I have been train	ismitted by the International Bureau					
		making such amendments has NOT expired.	}			
d. X have not been	made and will not be made.	-				
A translation of the am	endments to the claims under PCT	Article 19 (35 U.S.C. 371(c)(3)).				
An oath or declaration	of the inventor(s) (35 U.S.C. 371(c	:)(4)).				
A translation of the an (35 U.S.C. 371(c)(5)).	The state of the antickes to the international remaindry examination kenon under PL   Afficie 56					
Items 11. to 16. below concert	document(s) or information incli	uded:				
	sure Statement under 37 CFR 1.97	and 1.98,				
12. An assignment docum	ent for recording. A separate cover	sheet in compliance with 37 CFR 3.28 and 3.31 is mo	cluded.			
13. X A FIRST preliminary a	mendment					
A SECOND or SUBSE	QUENT preliminary amendment.					
14. A substitute specificati	on.					
15. A change of power of	attorney and/or address letter.					
16. X Other items or informa	tion:					
Courtesy copy o	f published PCT applicat	ion				
			1			
	-					
		·				

TOWNSEND&TOWNSEND

430 Rec'd PCT/PTO 2 7 SEP 2000

U.S. APPLIATOR WO. GEDAMA Sec 37 CFR LST - 7 7 DETERNATIONAL APPLICATION NO			ATTORNEYS DOCK	TNUMBER	
	19/64/327	PCT/GB99/00968		CALCULATIONS	PTO USE ONLY
17. X The following fees are submitted:  BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):  Neither international preliminary examination fee (37 CFR 1.482)  nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO  and International Search Report not prepared by the EPO or JPO					
International preliminary examination fee (37 CFR 1.482) not paid to (ISPTO) but International Search Report prepared by the EPO or IPO \$840.00					
International p	preliminary examination fee	(37 CFR 1.482) not paid to USP			
International preliminary examination fee paid to USPTO (37 CFR 1.482)					
but all claims did not satisfy provisions of PCT Article 33(1)-(4)			2)		
	_	RIATE BASIC FEE AM		\$ 840	
	0.00 for furnishing the oath earliest claimed priority dat		30	s	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total claims	14 -20 =		X \$18.00	\$	
Independent claims			X \$78.00	\$	
MULTIPLE DEP	ENDENT CLAIM(S) (if applic	· · · · · · · · · · · · · · · · · · ·	+ \$260.00	<u>s</u>	
1,1	TOTAL (	OF ABOVE CALCULAT	TONS =	s 840	
Reduction of 1/2 for filing by small entity, if applicable. A Small Entity Statement must also by filed (Note 37 CFR 1.9, 1.27, 1.28).			\$		
e the state of the		SUBT	OTAL =	\$	
Processing fee of \$130.00 for furnishing the English translation later than 20 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$	
TOTAL NATIONAL FEE =			\$ 840	-	
Fee fer recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property			S		
TOTAL FEES ENCLOSED =			_\$ 840		
			Amount to be:	S	
18 18 18 18 18 18 18 18 18 18 18 18 18 1				refunded charged	S
	_			<u> </u>	
a. A check in the amount of \$ to cover the above fees is enclosed,					
	charge my Deposit Account icate copy of this sheet is en	No. <u>20-1430</u> in the closed.	amount of \$ <u>84</u>	to cov	ver the above fees.
c. XX The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. 20-1430. A duplicate copy of this sheet is enclosed.					
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR s. 1.137(a) or (b)) must be filed and granted to restore the application to pending status.					
SEND ALL CORRESPONDENCE TO:					)
Joe L <b>i</b> ebes	schuetz			. Well-freeze	<u> </u>
	and Townsend and Cre		SIGNAT	IRE:	
	cadero Center, 8th f Lsco, CA 94111	1.	Jc	e Liebeschuetz	
Sen Etanci	LOCU, CA 74111		NAME	,	
				37,505	
			REGISTA	LATION NUMBER	

1

5

of:

Attorney Docket No.: 80359-000300US

## 430 Rec'd PCT/PTO 2 7 SEP 2000

#### IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re U.S. National Phase of

PCT/GB99/00968 of:

William Osmond Charles Michael Cookson et al.

Application No.: Unassigned

Filed: Herewith

For: POLYMORPHISM I: LINKAGE

OF ASTHMA TO A LOCUS ON

CHROMOSOME 2

PRELIMINARY AMENDMENT

Box PCT Assistant Commissioner for Patents Washington, D.C. 20231

Sir:

Prior to examination of the above-referenced application, please enter the following amendments and remarks.

#### IN THE CLAIMS:

Please amend claims 7, 8, 12 and 14 as follows. All pending unamended claims have been reproduced in small type for ease of reference.

- 1. (Amended under Article 34) A method for diagnosing an individual as being asthmatic, or as having a predisposition to asthma, which method comprises demonstrating in a nucleic acid sample taken from an individual the presence or absence of an allele which is associated with asthma, wherein the allele is situated at a locus in a region of chromosome 2 of up to 1 megabase in length, which region contains the locus D2S308.
  - 2. (As filed) The method according to claim 1, wherein the method comprises the steps
    - (i) Providing a suitable tissue sample from the individual;
    - (ii) Preparing from the tissue sample a nucleic acid sample
    - (iii) Analyzing the nucleic acid sample for the presence or absence of the allele.

2

1

1

2

1

2

3

1

2

1

2

1

2

3

1

2

3

1

2

1

2

3

1

2

William Osmond Charles Michael Cookson, et al. Application No.: Not yet assigned Page 2

PATENT

- (As filed) The method according to claim 2, wherein prior to analysis, the locus at which the allele is situated is amplified.
  - 4. (As filed) The method according to claim 3, wherein the amplification is by the PCR.
- 5. (As filed) The method according to any one of claims 1 to 4, wherein the locus at which the allele is situated comprises microsatellite repeats of variable lengths.
- 6. (As filed) The method according to claim 5, wherein amplification is performed using a pair of primers each of which hybridizes under suitably stringent conditions to a region either side of the microsatellite repeats.
- 7. (Amended) The method according to [any one of] claim[s] 1 [to 6], wherein the allele for identification is D2S308\*3.
- 8. (Amended) The method according to [any one of] claim[s] 3 [to 7], wherein the analysis is carried out by size separation of amplification products.
- (As filed) The method according to claim 6, wherein the primers in the pair of primers comprise the oligonucleotide sequences identified by SEQ ID NO: 1 and SEQ ID NO: 2 or substantially similar sequences.
- (As filed) A pair of oligonucleotide primers for amplification of an allele which is 10. associated with asthma, which allele is situated at a locus in a region of chromosome 2 of up to 1 megabase in length, which region contains the locus D2S308.
- (As filed) The pair of oligonucleotide primers according to claim 10, one of which is 11. labeled with a detectable marker.
- (Amended) The pair of oligonucleotides according to claim 10 [or claim 12. 11], capable of hybridizing under suitably stringent conditions to a region either side of a region of microsatellite repeats at D2S308.
- 13. (As filed) The pair of oligonucleotide primers according to claim 12, comprising the oligonucleotide sequences identified by SEQ ID NO: 1 and SEQ ID NO: 2 or substantially similar sequences.

William Osmond Charles Michael Cookson, et al. Application No.: Not yet assigned

Page 3

1

2

(Amended) An assay kit which comprises the pair of oligonucleotide 14. primers according to [any one of] claim[s] 10 [to 13]..

#### <u>REMARKS</u>

Amendments to claims 7, 8, 12 and 14 have been made to remove multiple claim dependency.

#### **CONCLUSION**

If the Examiner believes a telephone conference would expedite prosecution of this application, please telephone the undersigned at 650-326-2400.

Respectfully submitted,

**PATENT** 

Joe Liebeschuetz Reg. No. 37,505

TOWNSEND and TOWNSEND and CREW LLP Two Embarcadero Center,  $8^{\rm th}$  Floor

San Francisco, California 94111-3834

Telephone: (650) 326-2400 Facsimile: (650) 326-2422

JOL:dms PA 3099597 v1

10

15

20

25

# 0 9 / 6 4 7 3 2 9 430 Rec'd PCT/PTO 2 7 SEP 2000

#### POLYMORPHISM I : LINKAGE OF ASTHMA TO A LOCUS ON CHROMOSOME 2

This invention is concerned with methods for the diagnosis of asthma and with materials and methods relating thereto.

Asthma is a disease which is becoming more prevalent and is the most common disease of childhood (1). Most asthma in children and young adults is initiated by IgE mediated allergy (atopy) to inhaled allergens such as house dust mite and cat dander. However, not all asthmatics are atopic, and most atopic individuals do not have asthma, so that factors in addition to atopy are necessary to induce the disease (2,3). Asthma is strongly familial, and is due to the interaction between genetic and environmental factors. The genetic factors are thought to be variants of normal genes ("polymorphisms") which alter their function to predispose to asthma.

Asthma may be identified by recurrent wheeze and intermittent air flow limitation. An asthmatic tendency may be quantified by the measurement of bronchial hyper-responsiveness in which an individual's dose-response curve to a broncho-constrictor such as histamine or methacholine is constructed. The curve is commonly summarised by the dose which results in a 20% fall in air flow (PD20) or the slope of the curve between the initial air flow measurement and the last dose given (slope).

In the atopic response, IgE is produced by B-cells in response to allergen stimulation. These antibodies coat mast cells by binding to the high affinity receptor for IgE (FcɛRI). When a multivalent allergen binds to an IgE-coated mast cell, the cross-linking of adjacent IgEs by allergen initiates a series of cellular events leading to the destabilisation of the cell membrane

20

25

and release of inflammatory mediators. This results in mucosal inflammation, wheezing, coughing, sneezing and nasal blockage.

Atopy can be diagnosed by (i) a positive skin prick test in response to a common allergen; (ii) detecting the presence of specific serum IgE for allergen; or (iii) by detecting elevation of total serum IgE.

Genetic factors underlying a disease may be identified through localisation to particular chromosomal regions by genetic linkage. Genetic linkage is established by the study of families. It relies on matching the inheritance of disease with genetic polymorphisms of known localisation (known as "genetic markers"). In a complex disease such as asthma, genetic linkage will typically localise genes to within 10 - 20 Megabases (Mb) of DNA. A region of this size may contain 350 - 700 genes, and will be too large to permit immediate identification of the disease-causing gene.

Closer localisation of disease-causing genes may be accomplished by the detection of associations between particular alleles and the disease phenotype. Over short segments of DNA, distinctive alleles of the individual polymorphisms will show non-random association with alleles of neighbouring polymorphisms. This phenomenon, known as "linkage disequilibrium" occurs over 50-500 Kilobases (Kb) of DNA. Linkage disequilibrium may be detected by the study of individuals as well as by the study of families.

Disease-causing alleles will be in linkage disequilibrium with non-functional polymorphisms from the same chromosomal segment. It is therefore possible to detect allelic association with disease from particular chromosomal segments, without identifying the exact polymorphism and gene underlying the disease state.

The detection of allelic association may therefore give information as to disease susceptibility in a particular individual. Furthermore,

10

15

20

25

allelic association is indicative of a disease-causing gene being present within 500 Kb of DNA in either direction from the allele (i.e. 1 Mb in total). Such a region may contain only 30 genes, within which the identification of the disease-causing gene is possible.

The presence of linkage disequilibrium also means that other polymorphisms may be anticipated to associate with disease, and that these additional polymorphisms will also be diagnostic of disease susceptibility in particular individuals.

Genetic associations with atopy have been demonstrated. WO 95/05481 discloses that variants of the gene encoding the  $\beta$ -subunit of the high-affinity receptor for IgE (Fc $\epsilon$ RI $\beta$ ) are associated with atopy. It teaches a method for diagnosing atopy which is based upon the demonstration of the presence or absence of one of two variants in a specific portion of the DNA sequence of the gene encoding Fc $\epsilon$ RI $\beta$ , located near the commencement of exon 6 of the Fc $\epsilon$ RI $\beta$  gene on chromosome 11. A further variant has also been found in which the unusual variant sequence is in the coding sequence for the C-terminal cytoplasmic tail of Fc $\epsilon$ RI $\beta$  (4).

Tumour Necrosis Factor (TNF) is a pro-inflammatory cytokine that is found in increased concentration in asthmatic airways (5). We have previously shown that polymorphisms within the TNF gene are associated with an increased risk of asthma (6).

The known polymorphisms do not account for all of the genetic factors which predispose to asthma. In particular, asthma is not necessarily an atopic disease. Identification of further genetic polymorphisms linked to asthma will allow the identification of children at risk of asthma before the disease has developed (for example immediately after birth), with the potential for prevention of disease. The presence of particular polymorphisms may predict the clinical course of disease (e.g. severe as opposed to mild) or the

response to particular treatments. This diagnostic information will be of use to the health care, pharmaceutical and insurance industries.

It has now been discovered that a genetic polymorphism known as D2S308\*3 near the IL1 gene cluster on chromosome 2 is associated with asthma and can be used as a diagnostic tool.

Interleukin 1 (IL1) is another pro-inflammatory cytokine which is increased in asthmatic airways (5). IL1 is part of a gene family, localised on the long arm of chromosome 2 (7). Not all members of the gene family have been identified. The members of the family may share similar functions.

The invention therefore provides a method for diagnosing an individual as being asthmatic, or as having a predisposition to asthma, which method comprises demonstrating in the individual the presence or absence of an allele which is associated with asthma, wherein the allele is situated at a locus in a region of chromosome 2 of up to 1 megabase in length, which region contains the locus D2S308.

The 1 Mb region of chromosome 2 in which the D2S308 locus is situated flanks the D2S308 locus. Thus, the specific allele D2S308\*3, or other unusual polymorphisms in the region which are associated with asthma, may be the subject of identification in the method according to the invention. Equally, two or more such alleles may be the subject of identification.

Current diagnostic methods involving detection at the nucleic acid level normally comprise the steps of:

- (i) obtaining a suitable tissue sample from the individual;
- (ii) preparing from the tissue sample a nucleic acid sample;
- (iii) analysing the nucleic acid sample for the presence or absence of the relevant nucleic acid sequence, such as a specific allele.

10

20

15

25

10

15

20

25

Preferably, an amplification step is performed prior to the analysis, such that the locus at which the allele is situated is amplified. A preferred amplification technique is the PCR, although any suitable method of nucleic acid amplification may be employed.

In further aspects, the invention provides a pair of oligonucleotide primers for amplification of an allele which is associated with asthma, which allele is situated at a locus in a region of chromosome 2 of up to 1 megabase in length, which region contains the locus D2S308; and an assay kit comprising the pair of oligonucleotide primers.

The specific allele for identification may take the form of microsatellite repeats, which are nucleotide sequences containing short, repeated nucleotide motifs, usually a dinucleotide or a trinucleotide motif. A pair of primers which hybridize under suitably stringent conditions, to sequences at a position on either side of the microsatellite repeats, may be used to amplify the microsatellite repeats by PCR. Differences in the number of repeats are recognised by size differences in the PCR products. An allele which has a specified number of repeats and therefore a known size can thus be identified. D2S308\*3 is one such allele.

The primers employed in the method comprise nucleic acid sequences which are complementary to, or substantially complementary to unique sequences either side of the microsatellite repeats, such that only the relevant polymorphic region of the genome is amplified. The conditions under which the amplification is performed are gauged such that specific hybridization of the primers to the flanking sequences occurs and non-specific hybridization is avoided. The hybridization conditions are suitably stringent for that purpose. Standard techniques can be used to identify an appropriate set of reaction conditions.

Typically, the PCR products are detected by means of a detectable label attached to one of the PCR primers. Alternatively another form of labeling may be used such as a labeled sequence specific probe which hybridizes to the amplified sequences. The label may be a fluorescent or other label. The PCR products are subjected to size determination, typically involving size-separation for example by gel electrophoresis, and the presence or absence of the allele of interest is determined.

It will be evident that the invention is not limited with regard to the manner in which the presence or absence of the allele of interest is determined. The labeling, detection, separation or any other aspect of the method as described here may be replaced by other suitable known techniques and reagents.

The allele for identification may be an allele other than D2S308\*3 which is in linkage disequilibrium with D2S308\*3 and is associated with asthma. This includes alleles of both functional and non-functional polymorphisms. Functional polymorphisms include polymorphisms within genes, usually within coding sequences of genes. Non-functional polymorphisms are polymorphisms which do not themselves cause the disease.

This invention will now be further described in the Examples section which follows. The Examples are intended to be illustrative and do not limit the scope of the invention in any way.

#### **EXAMPLES**

Description of Laboratory Testing

#### Subjects

Three panels of subjects have been studied.

Panel A consisted of 80 nuclear families sub-selected from an

Australian population sample of 230 families (8). The panel contained a total of 203 offspring forming 172 sib-pairs. 12% of the children were asthmatic.

Panel B consisted of 77 nuclear and extended families recruited from asthma and allergy clinics in the United Kingdom (9). These families contained 215 offspring (268 sib-pairs) of which 56% were asthmatic.

Panel C consisted of 87 nuclear families recruited through a , child attending an asthma clinic in the Oxford region. The families contained 216 offspring (148 sibling pairs), of whom 44% were asthmatic.

#### 10 Phenotyping

5

15

25

The subjects were administered a modified British MRC respiratory questionnaire. "Asthma" was defined as a positive answer to the questions "Have you ever had an attack of asthma?" and "If yes, has this happened on more than one occasion?". "Wheeze" was defined as a positive answer to the question "Has your chest ever sounded wheezing or whistling?" and "If yes, has this happened on more than one occasion?". The total serum IgE was measured in all children. Skin tests to house dust mite and grass pollen were carried out.

#### 20 Genotyping

The microsatellite markers D2S340, D2S160, IL1A, D2S308 and D2S121 were typed by semi-automated fluorescent methods, as described previously (8). These markers cover a distance of 12 centiMorgans (approximately 12 million base pairs of DNA), containing the IL1 cluster of genes.

The polymerase chain reaction (PCR) primer sequences for the markers were as follows:

D2S308 5' GCC AGG TGG GTC TGG A (SEQ ID NO: 1)

UD2S308	5' TGG TTT TTT CAA GGG ATT TTC T (SEQ ID NO: 2)
D2S340	5' CTG GAA CCC CAG TAG C (SEQ ID NO: 3)
UD2S340	5' CTT TGC CCA GTT TTG C (SEQ ID NO: 4)
D2S160	5' TGT ACC TAA GCC CAC CCT TTA GAG C (SEQ ID NO: 5)
UD2S160	5' TGG CCT CCA GAA ACC TCC AA (SEQ ID NO: 6)
IL1A	5' GCC TAG TGA GTG TGG AAG ACA TTG (SEQ ID NO: 7)
UIL1A	5' CAG CAC TGG TTG GTC TTC ATC TTG (SEQ ID NO: 8)
D2S121	5' GCT GAT ATT CTG GTG GGA AA (SEQ ID NO: 9)
UD2S121	5' GGC AAG AGC AAA ACT CTGTC (SEQ ID NO: 10)

The polymerase chain reaction conditions were as follows: The reaction volumes were  $10\mu l$ , containing 50ng of genomic DNA, 200mM dNTPs, I x NH4+ buffer, 50ng oligonucleotide primers (forward labelled fluorescently), 0.5 to 3.0mM MgCl<sub>2</sub> and 0.2U Taq polymerase. Cycling conditions were 1 min at 95°C, 1min at 55°C and 45s at 72°C; 28 cycles were used. PCRs were performed on an Hybaid Omnigene thermal cycler.

Electrophoresis and allele scoring were as follows:

PCR products were mixed with a size standard (GS350 TAM) in loading buffer (80% (v/v) formamide, 20% (v/v) 50mM EDTA. 0.1% (w/v) blue dextran).

Samples were denatured at 95°C for 4min immediately prior to loading onto a 6% polyacrylamide gel and were electrophoresed at 800v for 6h on an Applied Biosystems (ABI) 373 DNA sequencer. Allele sizes were assigned using the ABI GENESCAN and ABI GENOTYPER software.

### Linkage and Association Analysis

Linkage was tested against asthma and the associated

10

15

20

phenotypes of wheeze, the skin test index, the total serum IgE and the residual IgE (the proportion of the total IgE not attributable to specific allergy)(8,10). Linkage was assessed by standard sib-pair methods as described (8).

Association was tested against the phenotype of asthma by the Transmission Disequilibrium Test.

#### Results

5

10

20

Genetic linkage to asthma-associated phenotypes was detected in Panel A and Panel B subjects as follows (genetic linkage to these phenotypes was not detected in Panel C subjects):

		Skin Te	st Index	Residu	ial IgE	Whe	eeze
	θ	Panel A	Panel B	Panel A	Panel B	Panel A	Panel B
D2S340	0.04	-	-	-	-	p<0.01	-
D2S160	0.02	-	p<0.005	p<0.01**	-	· p<0.01	p<0.05
IL1A	0.05	-	-	-		p<0.0005	-
D2S308	0.01	-	-	-	-	p<0.005	-
D2S121		-	-	-	-	p<0.01	-

<sup>\*</sup>recombination fraction: 0.01 approximates 1 million base pairs

Genetic linkage between the marker D2S160 and asthma was detected at p=0.09 in Panel A, p=0.01 in Panel B, and p=0.12 in Panel C.

The combined significance of linkage was p=0.008.

These results indicate that a gene influencing asthma lies within the D2S340 - D2S121 interval. Based on the current linkage information, the precise localisation of the putative asthma gene within the interval is uncertain.

<sup>15 \*\*</sup>p<0.00001 in maternal meioses

#### Association with D2S308 allele 3

Each of the markers was then tested for association with asthma phenotype. Association was seen in each panel for allele 3 of D2S308 (D2S308\*3). This allele is 228 base pairs in size, using the primers described above. (Other primers can be designed and their amplification product size determined for D2S308\*3, using known sequence in formation (11).) The results of TDT testing were as follows:

10		Transmitted	Not Transmitted	р
	Panel A	14	4	p=0.015
	Panel B	55	29	p=0.003
	Panel C	49	32	p=0.037
	Combined	118	65	p=0.00009

15

20

5

The results indicate that D2S308\*3, near the Interleukin 1 gene cluster on chromosome 2, shows a strong reproducible association with asthma in three diverse panels of subjects. It may therefore be inferred that a gene influencing asthma is present within 500 kilobases in either direction of D2S308.

#### REFERENCES

Strachan, D.P., Anderson, H.R., Limb, E.S., O'Neill, A. and Wells, N. (1994). A national survey of asthma prevalence, severity, and treatment in Great Britain. *Arch. Dis. Child.*, 70, 174-178.

15

25

- 2. Fraser, R.S., Paré, J.A.P., Fraser, R.G. and Paré, P.D., eds. (1994) Synopsis of Diseases of the Chest. WB Saunders Company, Philadelphia: 635-53.
- 3. Djukanovic, R., Roche, W.R., Wilson, J.W., Beasley, C.R., Twentyman, O.P., Howarth, R.H. and Holgate S.T. (1990) Mucosal inflammation in asthma. *Am. Rev. Respir. Dis.*, **142**, 434-457.
  - 4. Hill, M.R. and Cookson, W.O.C.M. (1996). A new variant of the β/subunit of the high-affinity receptor for Immunoglobulin E (FcεRI-β E237G): Associations of measures of atopy and bronchial hyper-responsiveness. *Hu. Mol. Gen.* 5, 959-962.
  - 5. Broide, D.H., Lotz, M., Cuomo, A.J., Coburn, D.A., Federman, E.C. and Wassserman, S.I. (1992) Cytokines in symptomatic asthma airways. *J. Allergy Clin. Immunol.* **89**, 958-967.
  - 6. Moffatt M.P, Cookson WOCM. Tumour Necrosis Factor Haplotypes. and Asthma. Hum Molec Genet 1997; 6: 551-554.
  - 7. Bailly S, di Giovine FS, Blakemore A.I.F, Duff G.W. Genetic polymorphism of human interleukin-1α. Eur J Immunol 1993; **23**: 1240-1245.
  - 8. Daniels S.E, Bhattacharyya S, James A, Leaves N.I, Young A, Hill M.R, Faux J.A, Ryan G.F, le Söuef P.N, Lathrop G.M, Musk A.W, Cookson
- 20 W.O.C.M. A genome-wide search for quantitative trait loci underlying asthma. Nature 1996; 383; 247-250.
  - 9. Sandford A.J, Shirakawa T, Moffatt M.F, Daniels S.E, Ra C, Faux J.A, Young R.P, Nakamura Y, Lathrop G.M, Cookson W.O.C.M Cookson, Hopkin J.M, Localisation of atopy and the β subunit of FcεRI receptor for IgE on chromosome 11g. Lancet 1993; 341: 332-334.
  - 10. Dizier M.H, Hill M, James A, Faux J, Ryan G, le Söuef P, Lathrop M, Musk A.W, Demenais F, Cookson W. Detection of a recessive major gene for

gang gingergene is to straig thing gives given by the straig ends gives give gang gives give a straight of the 
high IgE levels acting independently of specific responses to allergens. Genet Epidemiol 1995; **12**: 93-105.

11. Dib S. et al. A comprehensive map of the human genome based on 5264 microsatellites. Nature 1996; **380**: 152-154.

## CLAIMS AMENDED UNDER AZE PCT.

#### **CLAIMS**

- 1. A method for diagnosing an individual as being asthmatic, or as having a predisposition to asthma, which method comprises demonstrating in a nucleic acid sample taken from an individual the presence or absence of an allele which is associated with asthma, wherein the allele is situated at a locus in a region of chromosome 2 of up to 1 megabase in length, which region contains the locus D2S308.
- 2. The method according to claim 1, wherein the method comprises the steps of:
  - (i) providing a suitable tissue sample from the individual;
  - (ii) preparing from the tissue sample a nucleic acid sample;
  - (iii) analysing the nucleic acid sample for the presence or absence of the allele.
- 3. The method according to claim 2, wherein prior to analysis, the locus at which the allele is situated is amplified.
- 4. The method according to claim 3, wherein the amplification is by the PCR.
- 5. The method according to any one of claims 1 to 4, wherein the locus at which the allele is situated comprises microsatellite repeats of variable lengths.
- 6. The method according to claim 5, wherein amplification is performed using a pair of primers each of which hybridizes under suitably stringent conditions to a region either side of the microsatellite repeats.
- 7. The method according to any one of claims 1 to 6, wherein the allele for identification is D2S308\*3.

- 8. The method according to any one of claims 3 to 7, wherein the analysis is carried out by size separation of amplification products.
- The method according to claim 6, wherein the primers in the pair of primers comprise the oligonucleotide sequences identified by SEQ ID NO: 1 and SEQ ID NO: 2 or substantially similar sequences.
- 10. A pair of oligonucleotide primers for amplification of an allele which is associated with asthma, which allele is situated at a locus in a region of chromosome 2 of up to 1 megabase in length, which region contains the locus D2S308.
- 11. The pair of oligonucleotide primers according to claim 10, one of which is labelled with a detectable marker.
- 12. The pair of oligonucleotides according to claim 10 or claim 11, capable of hybridising under suitably stringent conditions to a region either side of a region of microsatellite repeats at D2S308.
- 13. The pair of oligonucleotide primers according to claim 12, comprising the oligonucleotide sequences identified by SEQ ID NO: 1 and SEQ ID NO: 2 or substantially similar sequences.
- 14. An assay kit which comprises the pair of oligonucleotide primers according to any one of claims 10 to 13.

Rev. 11-3/98

Effective Murch 1998

#### DECLARATION AND POWER OF ATTORNEY FOR U.S. PATENT APPLICATION

(X) Original () Supplemental () Substitute () PCT () DESIGN

As a below named inventor, I hereby declare that: my residence, post office address and citizenship are as stated below next to my name; that I verily believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

#### Title: POLYMORPHISM I: LINKAGE OF ASTHMA TO A LOCUS ON CHROMOSOME 2

of which is described and claimed in:

() the attached specification, or

(X) the specification in application Serial No. <u>09/647,329</u>, filed <u>September 27, 2000</u>, and with amendments through (if applicable), or

() the specification in International Application No. , filed , and as amended on (if applicable).

I hereby state that I have reviewed and understand the content of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge my duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, §1.56.

I hereby claim priority benefits under Title 35, United States Code, §119 (and §172 if this application is for a Design) of any is application application for patent or inventor's ertificate listed below and have also identified below any application for patent or inventor's ertificate having a filing date before that of the application on which priority is claimed:

COUNTRY	APPLICATION NO.	DATE OF FILING	PRIORITY CLAIMED
Great Britain	9806652.5	March 27, 1998	Yes

I hereby claim the benefit under Title 35, United States Code §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code §112, I acknowledge the duty to disclose information material to patentability as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application.

APPLICATION SERIAL NO.	U.S. FILING DATE	STATUS: PATENTED, PENDING, ABANDONED

And I hereby appoint Michael R. Davis, Reg. No. 25,134; Matthew M. Jacob, Reg. No. 25,154; Jeffrey Nohon, Reg. No. 25,408; Warren M. Cheek, Jr., Reg. No. 33,367; Nils Pedersen, Reg. No. 33,145; and Charles R. Watts, Reg. No. 33,142, who together constitute the firm of WENDEROTH, LIND & PONACK, L.L.P., jointly and severally, attorneys to prosecute this application and to transact all business in the U.S. Patent and Trademark Office connected therewith.

I hereby authorize the U.S. attorneys named herein to accept and follow instructions from <u>Kilburn and Strode</u> as to any action to be taken in the U.S. Patent and Trademark Office regarding this application without direct communication between the U.S. attorneys and myself. In the event of a change in the persons from whom instructions may be taken, the U.S. attorneys named herein will be so notified by me.

Send Correspondence to

Direct Telephone Calls to:

WENDEROTH, LIND & PONACK, L.L.P. 2033 K Street, N.W., Suite 800 Washington, D.C. 20006

WENDEROTH, LIND & PONACK, L.L.P. Area Code (202) 721-8200

Direct Facsimile Messages to: Area Code (202) 721-8250

Full Name of First Inventor	FAMILY NAME FIRST CIVEN NAME SECOND GIVEN NAME  COOKSON William Osmond Charles Michael
Residence & Citizenship	CITY STATE OR COUNTRY COUNTRY OF CITIZENSHIP Oxford Great Britain Great Britain
Post Office Address	ADDRESS CITY STATEOR COUNTRY ZIP CODE 67 Hilltop Road Oxford Great Britain OX4 1PD
Full Name of Second Inventor	FAMILY NAME FIRST GIVEN NAME SECOND GIVEN NAME MOFFATT Miriam Fleur
Residence & Citizenship	CITY STATE OR COUNTRY COUNTRY OF CITIZENSHI  Langford Village, Bicester Great Britain Great Britain
Post Office Address	Address City STATE OR COUNTRY ZIP CODE 30 Lapwing Close Langford Village, Bicester Great Britain OX6 0XR
Full Name of _ Third Inventor	FAMILY NAME FIRST CIVEN NAME SECOND GIVEN NAME BHATTACHARYYA Sumit
Residence & Citizenship	Stapleford, Cambridgeshire Great Britain Great Britain
Post Office Address	ADDRESS CITY STATE OR COUNTRY ZIP COD 20 Bury Road Stapleford, Cambridgeshire Great Britain CB2 5BP
Full Name of Fourth Inventor	FAMILY NAME FIRST GIVEN NAME SECOND GIVEN NAME LEAVES Nicholas
Residence & Citizenship	Sandhills, Oxford Great Britain G
Post Office Address	ADDRESS CITY STATE OR COUNTRY ZIP CODE 47 Burdell Avenue Sandhills, Oxford Great Britain OX3 8EE
Full Name of Fifth Inventor	PAMILY NAME FIRST GIVEN NAME SECOND GIVEN NAME
Residence & Citizenship	CITY STATE OR COUNTRY COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS CITY STATE OR COUNTRY ZIP CODE
Full Name of Sixth Inventor	FAMILY NAME FIRST GIVEN NAME SECOND GIVEN NAME
Residence & Citizenship	CITY STATE OR COUNTRY COUNTRY OF CITIZENSHIP
Post Office Address	ADDRESS CITY STATE OR COUNTRY ZIP CODE

I further declare that all statements made herein of my own knowledge are true, and that all statements on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent/issuing thereon.

1st Inventor Littam Cooker	_ X Date 23.1.01	_
William Osmond Charles Midfinel COOKSON  2nd Inventor	X Date 23.1.01	×
Miriam Flour MOFFATT  3rd Inventor Alexandrage	X Date 14-12-16-1	- ~
SuninBHATTACHARYYA	× <sub>Date</sub> 31.1.01	-^
Ath Inventor / OC// GC FY / O. 1	Date 3( ) ( )	- /-
5th Inventor	Date	-
6th Inventor	Date	_

The above application may be more particularly identified as follows:

U.S. Application Serial No. 09/647,329 Filling Date September 27, 2000

Applicant Reference Number P30627US/KVC/PS/CF Atty Docket No. 2000-1718

01865-272412

Title of Invention POLYMORPHISM I: LINKAGE OF ASTHMA TO A LOCUS ON CHROMOSOME 2